arXiv:1606.02040v1 [math.FA] 7 Jun 2016

COARSE LIPSCHITZ EMBEDDINGS OF JAMES SPACES

F. NETILLARD

Abstract. We prove that, for $1 < p \neq q < \infty$, there does not exist any coarse Lipschitz embedding between the two James spaces J_p and J_q, and that, for $1 < p < q < \infty$ and $1 < r < \infty$ such that $r \notin \{p, q\}$, J_r does not coarse Lipschitz embed into $J_p \oplus J_q$.

1. Introduction

Let (M, d) and (N, δ) be two metric spaces and $f : M \to N$.
The map f is said to be a *coarse Lipschitz embedding* if there exist θ, A, $B > 0$ such that

$$\forall\ x,\ y \in M \quad d(x, y) \geq \theta \Rightarrow Ad(x, y) \leq \delta(f(x), f(y)) \leq Bd(x, y).$$

Then we say that M *coarse Lipschitz embeds* into N.

R.C. James introduced in [5] a non-reflexive space defined by :

$$J = \left\{ x : \mathbb{N} \to \mathbb{R} \text{ s.t. } x(n) \to 0 \text{ and } ||x||_J = \sup_{p_1 < \ldots < p_n} \left(\sum_{i=1}^{n-1} |x(p_{i+1}) - x(p_i)|^2 \right)^{\frac{1}{2}} < \infty \right\}$$

We will use the following spaces (where $1 < p < \infty$), which are variants of J :

$$J_p = \left\{ x : \mathbb{N} \to \mathbb{R} \text{ s.t. } x(n) \to 0 \text{ and } ||x||_{J_p} = \sup_{p_1 < \ldots < p_n} \left(\sum_{i=1}^{n-1} |x(p_{i+1}) - x(p_i)|^p \right)^{\frac{1}{p}} < \infty \right\}$$

Like in the case of J, the codimension of J_p in J_p^{**} is 1.
In this respect, we precise that J_p^{**} can be seen as :

$$J_p^{**} = \left\{ x : \mathbb{N} \to \mathbb{R} \text{ s.t. } \sup_{p_1 < \ldots < p_n} \left(\sum_{i=1}^{n-1} |x(p_{i+1}) - x(p_i)|^p \right)^{\frac{1}{p}} < \infty \right\}$$

All those spaces were introduced in [11].

In 2007, N.J. Kalton and N.L. Randrianarivony [5] proved that, if $r \notin \{p_1,\ \ldots,\ p_n\}$ where $1 \leq p_1 < p_2 < \ldots < p_n < \infty$, then ℓ_r does not coarse Lipschitz embed into $\ell_{p_1} \oplus \ldots \oplus \ell_{p_n}$.
The aim of this article is to prove similar results for the J_p spaces. One of the main obstacles is the lack of reflexivity, which was crucial in Kalton-Randrianarivony's

2010 *Mathematics Subject Classification.* Primary 46B20; Secondary 46B80 .
Key words and phrases. Banach spaces, James spaces, coarse Lipschitz embeddings.

work. However, the James spaces have nice properties of asymptotic uniform smoothness and weak* asymptotic uniform convexity that we shall use (see [8] or [9] for the definitions). We shall not refer to these notions in our paper, but we will build concrete equivalent norms on J_p that will serve our purpose. Some compactness arguments will also be used to deal with the extra dimension in J_p^{**}.

This paper is organized as follows. In Section 2 we summarize the notation and terminology and we give the basic results. Section 3 contains the proof of the nonexistence of a coarse Lipschitz embedding between two James spaces J_p and J_q for $1 < p \neq q < \infty$. At the end of this last section, we show that, for $1 < p < q < \infty$ and $1 < r < \infty$ such that $r \notin \{p, q\}$, J_r does not coarse Lipschitz embed into $J_p \oplus J_q$.

2. Preliminaries

Notation 2.1. Let e_n defined by $e_n(k) = \delta_{n,k}$ for $k \in \mathbb{N}$. The sequence $(e_n)_{n=0}^{\infty}$ is a Schauder basis of J_p (where $p > 1$).
Moreover, the sequence $(e_n^*)_{n=0}^{\infty}$ of the coordinate functionals associated with $(e_n)_{n=0}^{\infty}$ is a Schauder basis of J_p^*.
When u and v in J_p have a consecutive and disjoint finite supports with respect to $(e_n)_{n=0}^{\infty}$, we will denote $u \prec v$.
Likewise, when u^* and v^* in J_p^* have a consecutive and disjoint finite supports with respect to $(e_n^*)_{n=0}^{\infty}$, we will denote $u^* \prec v^*$.

We start with the construction of an ad'hoc equivalent norm on J_p. We follow the construction given in [10] for J_2.

Lemma 2.2. *Let $x_1, \ldots, x_n$ in J_p such that their supports are consecutive and finite with respect to the basis $(e_n)_{n=0}^{\infty}$. Then*

$$||\sum_{i=1}^{n} x_i||_{J_p}^p \leq (2^p + 1) \sum_{i=1}^{n} ||x_i||_{J_p}^p.$$

Proof. For $x \in J_p$, we denote $supp(x) = \{n \in \mathbb{N},\ e_n^*(x) \neq 0\}$. Then we can find disjoint intervals in $\mathbb{N}$, $[\![p_i,\ p_i']\!]$, with $1 \leq i \leq n$ and $p_i' < p_{i+1}$, such that :
$\forall\ 1 \leq i \leq n,\ supp(x_i) \subset [\![p_i,\ p_i']\!]$ (for convenience, we fix $p_1 = 0$ et we denote $p_{n+1} = \infty$).
Let now $q_1 < ... < q_k$ be an arbitrary sequence in $\mathbb{N}$. We must show that

$$\sum_{j=1}^{k-1} |y(q_j) - y(q_{j+1})|^p \leq (2^p + 1) \sum_{i=1}^{n} ||x_i||_{J_p}^p,$$

where $y = \sum_{i=1}^{n} x_i$.
There exist an increasing sequence $(j_m)_{m=1}^{l}$ in $\{1, \ldots, k\}$ with $j_1 = 1$, and an increasing sequence $(i_m)_{m=1}^{l}$ in $\{1, \ldots, n\}$ such that, for any $1 \leq m \leq l$, $\{q_{j_m}, \ldots, q_{j_{m+1}} - 1\} \subset [\![p_{i_m}, p_{i_{m+1}}]\!]$. Therefore

$$\sum_{j=1}^{k-1} |y(q_j) - y(q_{j+1})|^p =$$

$$\sum_{j=1}^{j_2-1} |y(q_j)-y(q_{j+1})|^p + |y(q_{j_2-1})-y(q_{j_2})|^p + ... + \sum_{j=j_{l-1}}^{j_l-1} |y(q_j)-y(q_{j+1})|^p .$$

We have that

$$\forall\ 1 \le m \le l-1,\ \sum_{j=j_m}^{j_{m+1}-1} |y(q_j)-y(q_{j+1})|^p \le ||x_{i_m}||^p_{J_p}.$$

And for all $2 \le m \le l-1$,

$$|y(q_{j_m-1})-y(q_{j_m})|^p \le 2^{p-1}\,|y(q_{j_m-1})|^p + 2^{p-1}\,|y(q_{j_m})|^p \le 2^{p-1}||x_{i_m}||^p_{J_p} + 2^{p-1}||x_{i_{m+1}}||^p_{J_p}.$$

Then

$$\sum_{j=1}^{k-1}(y(q_j)-y(q_{j+1}))^p \le (2^{p-1}+1)||x_{i_1}||^p_{J_p} + (2^p+1)||x_{i_2}||^p_{J_p} + ... + (2^p+1)||x_{i_{l-1}}||^p_{J_p} + (2^{p-1}+1)||x_{i_l}||^p_{J_p}.$$

This concludes the proof of our Lemma. □

We now define a new norm on J_p^* as follows. Let q be the conjugate exponent of p. For $x^* \in J_p^*$, we set

$$|x^*|_{J_p^*} = \sup\Big\{ (\sum_{i=1}^{n} ||x_i^*||^q_{J_p^*})^{\frac{1}{q}}\ : x^* = x_1^* + ... + x_n^*\ et\ x_1^* \prec ... \prec x_n^* \Big\},$$

where $||.||_{J_p^*}$ denotes the dual norm of $||.||_{J_p}$. We can now state the following proposition.

Proposition 2.3. *The norm $|.|_{J_p^*}$ is the dual norm of an equivalent norm on J_p (that we shall denote $|.|_{J_p}$).*
Moreover, $|.|_{J_p^*}$ satisfies the following property: for any x^*, y^* in J_p^* such that $x^* \prec y^*$, we have that

$$|x^*+y^*|^q_{J_p^*} \ge |x^*|^q_{J_p^*} + |y^*|^q_{J_p^*}.$$

Proof. To show that $|.|_{J_p^*}$ is a norm, we only detail the proof of the triangle inequality : let $(x^*,\ y^*) \in \left(J_p^*\right)^2$ that we may assume with finite supports. Let now $u_1^* \prec u_2^* \prec ... \prec u_n^*$ in J_p^* such that

$$x^* + y^* = u_1^* + ... + u_n^*$$

.

We write, for $i \in [\![1,\ n]\!]$, $u_i^* = x_i^* + y_i^*$, where $x^* = \sum_{i=1}^{n} x_i^*$ and $y^* = \sum_{i=1}^{n} y_i^*$.
Thanks to the triangle inequality for $||.||_{J_p^*}$, we get:

$$\left(\sum_{i=1}^{n} ||u_i^*||^q_{J_p^*}\right)^{\frac{1}{q}} \le \left(\sum_{i=1}^{n} \left(||x_i^*||_{J_p^*} + ||y_i^*||_{J_p^*}\right)^q\right)^{\frac{1}{q}}.$$

It then follows from Minkowski's inequality that

$$\left(\sum_{i=1}^{n}||u_i^*||_{J_p^*}^q\right)^{\frac{1}{q}} \leq \left(\sum_{i=1}^{n}||x_i^*||_{J_p^*}^q\right)^{\frac{1}{q}} + \left(\sum_{i=1}^{n}||y_i^*||_{J_p^*}^q\right)^{\frac{1}{q}} \leq |x^*|_{J_p^*} + |y^*|_{J_p^*}.$$

We have shown that the triangle inequality is valid for $|.|_{J_p^*}$.

Next we show that there exists $c > 0$ (which will be detailed later) such that, for x_1^*, ..., x_n^* in J^* satisfying $x_1^* \prec ... \prec x_n^*$ with respect to the basis $(e_n^*)_{n=0}^{\infty}$:

$$||\sum_{i=1}^{n} x_i^*||_{J_p^*}^q \geq c\sum_{i=1}^{n}||x_i^*||_{J_p^*}^q. \tag{2.1}$$

So, let $x_1^* \prec ... \prec x_n^*$, with, for $i \in [\![1,\ n]\!]$, $supp(x_i^*) \subseteq [\![p_i,\ q_i]\!]$, where $q_i < p_{i+1}$ for $i \in [\![1,\ n-1]\!]$. Fix now $\varepsilon > 0$.
Since $(e_i)_{i=0}^{\infty}$ is a monotone basis:

$$\exists\ x_i \in J_p,\ \begin{cases} x_i^*(x_i) & \geq & ||x_i^*||_{J_p^*}^q - \varepsilon \\ ||x_i||_{J_p} & \leq & 2||x_i^*||_{J_p^*}^{q-1} \\ supp(x_i) & \subseteq & [\![p_i,\ q_i]\!] \end{cases}$$

Thanks to Lemma 2.2 : $\exists\ C > 0,\ ||\sum_{i=1}^{n} x_i||_{J_p}^p \leq C\sum_{i=1}^{n}||x_i||_{J_p}^p$ (where $C = 2^p + 1$).
Since $||.||_{J_p^*}$ is the dual norm of $||.||_{J_p}$, we have that

$$||\sum_{i=1}^{n} x_i^*||_{J_p^*} \geq \left(\sum_{i=1}^{n} x_i^*\right)\left(\sum_{i=1}^{n} x_i\right)\left(||\sum_{i=1}^{n} x_i||_{J_p}\right)^{-1}$$

and

$$||\sum_{i=1}^{n} x_i^*||_{J_p^*} \geq \left(\sum_{i=1}^{n} x_i^*(x_i)\right)\left(||\sum_{i=1}^{n} x_i||_{J_p}\right)^{-1} \geq \left(\sum_{i=1}^{n}||x_i^*||_{J_p^*}^q - \varepsilon\right)\left(C^{\frac{1}{p}}\left(\sum_{i=1}^{n}||x_i||_{J_p}^p\right)^{\frac{1}{p}}\right)^{-1}.$$

Moreover, $\left(\sum_{i=1}^{n}||x_i||_{J_p}^p\right)^{\frac{1}{p}} \leq 2\left(\sum_{i=1}^{n}||x_i^*||_{J_p^*}^{p(q-1)}\right)^{\frac{1}{p}} = 2\left(\sum_{i=1}^{n}||x_i^*||_{J_p^*}^q\right)^{\frac{1}{p}}$.
Letting ε tend to 0, we obtain :

$$||\sum_{i=1}^{n} x_i^*||_{J_p^*} \geq \frac{1}{2C^{\frac{1}{p}}}\left(\sum_{i=1}^{n}||x_i^*||_{J_p^*}^q\right)^{1-\frac{1}{p}} = \frac{1}{2C^{\frac{1}{p}}}\left(\sum_{i=1}^{n}||x_i^*||_{J_p^*}^q\right)^{\frac{1}{q}}.$$

So, we have established inequality (2.1) with $c = \dfrac{1}{2^q C^{\frac{q}{p}}} = \dfrac{1}{2^q C^{q-1}}$.
It follows easily that

$$||x^*||_{J_p^*} \leq |x^*|_{J_p^*} \leq c^{-1/q}||x^*||_{J_p^*}.$$

Moreover, $|.|_{J_p^*}$ is the dual norm of an equivalent norm on J_p. Indeed, it is clear that $|.|_{J_p^*}$ is $\sigma(J_p^*, J_p)$ lower semi-continuous.

Finally, it follows clearly from the definition of $|.|_{J_p^*}$ that for all x^*, y^* in J_p^* such that $x^* \prec y^*$, we have that

$$|x^* + y^*|_{J_p^*}^q \geq |x^*|_{J_p^*}^q + |y^*|_{J_p^*}^q.$$

□

Corollary 2.4. *The dual norm $|.|_{J_p^{**}}$ of $|.|_{J_p^*}$ satisfies the following property. For $x \in J_p$ with a finite support and $y \in J_p^{**}$ (not necessarily with finite support) such that $x \prec y$, we have*

$$|x + y|_{J_p^{**}}^p \leq |x|_{J_p^{**}}^p + |y|_{J_p^{**}}^p.$$

Proof. Let $x \in J_p$ which has a finite support and $y \in J_p^{**}$ such that $x \prec y$, with $supp(x) \subset [\![m,\ n]\!]$, $supp(y) \subset [\![m',\ +\infty)$ and $n < m'$. Fix $\varepsilon > 0$.
There exists $z^* \in J_p^*$ such that

$$|z^*|_{J_p^*} = |x + y|_{J_p^{**}}^{p-1} \quad \text{and} \quad z^*(x + y) \geq |x + y|_{J_p^{**}}^p - \varepsilon.$$

Moreover, we can write $z^* = x^* + y^*$, with $x^* \prec y^*$, $z^*(x) = x^*(x)$ and $z^*(y) = y^*(y)$. We deduce that $|x + y|_{J_p^{**}}^p \leq x^*(x) + y^*(y) + \varepsilon$. Then Hölder's inequality and Proposition 2.3 yield

$$|x+y|_{J_p^{**}}^p \leq (|x^*|_{J_p^*}^q + |y^*|_{J_p^*}^q)^{\frac{1}{q}}(|x|_{J_p^{**}}^p + |y|_{J_p^{**}}^p)^{\frac{1}{p}} + \varepsilon \leq (|x^* + y^*|_{J_p^*})(|x|_{J_p^{**}}^p + |y|_{J_p^{**}}^p)^{\frac{1}{p}} + \varepsilon.$$

Since $|z^*|_{J_p^*} = |x + y|_{J_p^{**}}^{p-1}$, we get

$$|x + y|_{J_p^{**}}^p \leq (|x + y|_{J_p^{**}}^{p-1})(|x|_{J_p^{**}}^p + |y|_{J_p^{**}}^p)^{\frac{1}{p}} + \varepsilon.$$

This finishes our proof. □

We now turn to the study of the coarse Lipschitz embeddings between James spaces. Let us first recall some notation.

Definition 2.5. Let (M, d) and (N, δ) be two metric spaces and $f : M \to N$ be a mapping. If (M, d) is unbounded, we define

$$\forall s > 0, \quad Lip_s(f) = \sup\left\{\frac{\delta((f(x), f(y))}{d(x, y)},\ d(x, y) \geq s\right\} \quad \text{and} \quad Lip_\infty(f) = \inf_{s>0} Lip_s(f).$$

Note that f is coarse Lipschitz if and only if $Lip_\infty(f) < \infty$.

We also recall a classical definition.

Definition 2.6. Given a metric space X, two points $x,\ y \in X$, and $\delta > 0$, the approximate metric midpoint set between x and y with error δ is the set :

$$Mid(x,\ y,\ \delta) = \left\{z \in X\ :\ \max\{d(x,\ z),\ d(y,\ z)\} \leq (1 + \delta)\frac{d(x,\ y)}{2}\right\}$$

The use of approximate metric midpoints in the study of nonlinear geometry is due to Enflo in an unpublished paper and has been used elsewhere, e.g. [2], [3] and [6]. The next proposition and its proof can be found for instance in [8] and [9].

Proposition 2.7. *Let X be a normed space and suppose M is a metric space. Let $f : X \to M$ be a coarse Lipschitz map. If $Lip_\infty(f) > 0$, then for any $t,\ \varepsilon > 0$ and any $0 < \delta < 1$, there exist $x,\ y \in X$ with $||x - y|| > t$ and*

$$f\left(Mid(x,\ y,\ \delta)\right) \subset Mid\left(f(x),\ f(y),\ (1+\varepsilon)\delta\right).$$

Let us now recall the definition of the metric graphs introduced in [8] that will be crucial in our proofs.

Notation 2.8. Let $\mathbb{M}$ be an infinite subset of $\mathbb{N}$ and $k \in \mathbb{N}$. We denote

$$G_k(\mathbb{M}) = \{\overline{n} = (n_1, ..., n_k),\ n_i \in \mathbb{M} \quad n_1 < ... < n_k\}.$$

Then we equip $G_k(\mathbb{M})$ with the distance $d(\overline{n}, \overline{m}) = |\{j,\ n_j \neq m_j\}|$.

We end these preliminaries by recalling Ramsey's theorem and one of its immediate corollaries (see [4] for instance).

Theorem 2.9. *Let $k,\ r \in \mathbb{N}$ and $f : G_k(\mathbb{N}) \to \{1, ..., r\}$ be any map. Then there exists an infinite subset $\mathbb{M}$ of $\mathbb{N}$ and $i \in \{1, ..., r\}$ such that, for every $\overline{n} \in G_k(\mathbb{M})$, $f(\overline{n}) = i$.*

Corollary 2.10. *Let (K, d) be a compact metric space, $k \in \mathbb{N}$ and $f : G_k(\mathbb{N}) \to K$. Then for every $\epsilon > 0$, there exist an infinite subset $\mathbb{M}$ of $\mathbb{N}$ such that for every $\overline{n}, \overline{m} \in G_k(\mathbb{M}),\ d(f(\overline{n}), f(\overline{m})) < \epsilon$.*

3. The main results

Our first Lemma gives a description of approximate metric midpoints in J_p that is analogous to situation in ℓ_p (see [8] or [9]). However, we need to use both the original and our new norm on J_p.

Lemma 3.1. *Let $1 < p < \infty$. We denote E_N the closed linear span of $\{e_i,\ i > N\}$. Let now $x,\ y \in J_p,\ \delta \in (0,\ 1)$, $u = \dfrac{x+y}{2}$ and $v = \dfrac{x-y}{2}$. Then*
(i) There exists $N \in \mathbb{N}$ such that:

$$u + \delta^{\frac{1}{p}} |v|_{J_p} B_{(E_N, |.|_{J_p})} \subset Mid_{|.|_{J_p}}(x,\ y,\ \delta).$$

(ii) There is a compact subset K of J_p such that:

$$Mid_{||.||_{J_p}}(x,\ y,\ \delta) \subset K + 2\delta^{\frac{1}{p}} ||v||_{J_p} B_{(J_p, ||.||_{J_p})}.$$

Proof. Fix $\lambda > 0$.
Let $N \in \mathbb{N}$ such that $|v - v_N|_{J_p} \leq \lambda |v|_{J_p}$ and $|v_N|^p_{J_p} \geq (1 + \lambda^p)^{-1} |v|^p_{J_p}$, where $v_N = \sum_{i=1}^N v(i) e_i$.

(i) Let now $z \in E_N$ so that $|z|^p_{J_p} \leq \delta |v|^p_{J_p}$. Then

$$|x - (u+z)|^p_{J_p} = |v - z|^p_{J_p} = |v - v_N + v_N - z|^p_{J_p}$$

It follows from the Corollary 2.4 that:

$$|x - (u+z)|^p_{J_p} \leq |v - v_N - z|^p_{J_p} + |v_N|^p_{J_p} \leq (|v - v_N|_{J_p} + |z|_{J_p})^p + |v|^p_{J_p}$$

Therefore
$$|x-(u+z)|_{J_p}^p \leq \big((1+\delta^{1/p})^p+1\big)|v|_{J_p}^p \leq (1+\delta)^p|v|_{J_p}^p,$$
if λ was chosen initially small enough.
We argue similarly to show that $|y-(u+z)|_{J_p} = |v+z|_{J_p} \leq (1+\delta)|v|_{J_p}$ and deduce that $u+z \in Mid(x,\ y,\ \delta)$.

(ii) Fix $\nu > 0$ and choose $N \in \mathbb{N}$ such that $\|v_N\|_{J_p}^p \geq (1-\nu^p)\|v\|^p$. We assume now that $u+z \in Mid_{||.||_{J_p}}(x,\ y,\ \delta)$ and write $z = z' + z''$ with $z' \in F_N = sp\{e_i, i \leq N\}$ and $z'' \in E_N$.
Since $||v-z||_{J_p}\ ,\ ||v+z||_{J_p} \leq (1+\delta)||v||_{J_p}$, we get, by convexity, that
$$||z'||_{J_p} \leq ||z||_{J_p} \leq (1+\delta)||v||_{J_p}.$$
Therefore, $u+z'$ belongs to the compact set $K = u + (1+\delta)||v||_{J_p}B_{(F_N,||.||_{J_p})}$.
Moreover, for any $(m,n) \in (\mathbb{N}^*)^2$, with $m > n$:
$$\max\{|v(n)-v(m)|^p,\ |z(n)-z(m)|^p\} \leq \frac{1}{2}(|(v(n)-z(n))-(v(m)-z(m))|^p + |(v(n)+z(n))-(v(m)+z(m))|^p).$$
Therefore
$$(1-\nu^p)||v||_{J_p}^p + ||z''||_{J_p}^p \leq ||v_N||_{J_p}^p + ||z''||_{J_p}^p \leq \frac{1}{2}(||v-z||_{J_p}^p + ||v+z||_{J_p}^p) \leq (1+\delta)^p||v||_{J_p}^p.$$
Then, if ν was chosen small enough, we get
$$||z''||_{J_p}^p \leq [(1+\delta)^p - (1-\nu^p)]||v||_{J_p}^p \leq 2^p\delta||v||_{J_p}^p.$$

□

Proposition 3.2. *Let $1 < p < q < \infty$ and $f : (J_q,\ |.|_{J_q}) \to (J_p,\ ||.||_{J_p})$ be a coarse Lipschitz embedding. Then, for any $t > 0$ and for any $\varepsilon > 0$, there exist $u \in J_q$, $\theta > t$, $N \in \mathbb{N}$ and K a compact subset of J_p such that*
$$f(u + \theta B_{(E_N,|.|_{J_q})}) \subset K + \varepsilon\theta B_{(J_p,||.||_{J_p})}.$$

Proof. If $Lip_\infty(f) = 0$, the conclusion is clear. So we assume that $Lip_\infty(f) > 0$.
We choose a small $\delta > 0$ (to be detailed later). Then we choose s large enough so that $Lip_s(f) \leq 2Lip_\infty(f)$.
Then, by Proposition 2.7,
$$\exists\ x,\ y \in J_q,\ |x-y|_{J_q} \geq s \text{ and } f(Mid_{|.|_{J_q}}(x,\ y,\ \delta)) \subset Mid_{||.||_{J_p}}(f(x),\ f(y),\ 2\delta).$$
Denote $u = \dfrac{x+y}{2}$, $v = \dfrac{x-y}{2}$ and $\theta = \delta^{\frac{1}{q}}|v|_{J_q}$. By Lemma 3.1, there exists $N \in \mathbb{N}$ such that $u+\theta B_{(E_N,|.|_{J_q})} \subset Mid_{|.|_{J_q}}(x,\ y,\ \delta)$ and there exists a compact subset K of J_p so that $Mid_{||.||_{J_p}}(f(x),\ f(y),\ 2\delta) \subset K + 2(2\delta)^{\frac{1}{p}}||f(x)-f(y)||_{J_p}B_{(J_p,||.||_{J_p})}$. But :
$$2(2\delta)^{\frac{1}{p}}||f(x)-f(y)||_{J_p} \leq 4Lip_\infty(f)(2\delta)^{\frac{1}{p}}|x-y|_{J_q} \leq 8Lip_\infty(f)2^{\frac{1}{p}}\delta^{\frac{1}{p}-\frac{1}{q}}\theta \leq \varepsilon\theta,$$
if δ was chosen initially small enough.

Then an appropriate choice of a large s will ensure that $\theta \geq \frac{1}{2}\delta^{\frac{1}{q}} s > t$. This finishes the proof. □

Corollary 3.3. *Let $1 < p < q < \infty$.*
Then J_q does not coarse Lipschitz embed into J_p.

Proof. We proceed by contradiction and suppose that there exists a coarse Lipschitz embedding $f : (J_q, |.|_{J_q}) \to (J_p, ||.||_{J_p})$.
With the notation of the previous proposition, we can find a sequence $(u_n)_{n=1}^{\infty}$ in $u + \theta B_{(E_N, |.|_{J_q})}$, such that $|u_n - u_m|_{J_q} \geq \theta$ for $n \neq m$. Then $f(u_n) = k_n + \varepsilon\theta v_n$, with $k_n \in K$ et $v_n \in B_{(J_p, ||.||_{J_p})}$. Since K is compact, by extracting a subsequence, we may assume that $||f(u_n) - f(u_m)||_{J_p} \leq 3\varepsilon\theta$.
Since ε can be chosen arbitrarily small and θ arbitrarily large, this yields a contradiction. □

In order to treat the coarse Lipschitz embeddability in the other direction, we shall use the Kalton-Randrianarivony graphs and some special subsets of them that we introduce now.

Definition 3.4. Let $\overline{n}$, $\overline{m} \in G_k(\mathbb{M})$ (where $\mathbb{M}$ is an infinite subset of $\mathbb{N}$).
We say that $(\overline{n}, \overline{m}) \in I_k(\mathbb{M})$ if $n_1 < m_1 < n_2 < m_2 < ... < n_k < m_k$.

Proposition 3.5. *Let $\varepsilon > 0$ and $f : G_k(\mathbb{N}) \to (J_p^{**}, |\ |_{J_p^{**}})$ be a Lipschitz map.*
Then, for any infinite subset $\mathbb{M}$ of $\mathbb{N}$, there exists $(\overline{n}, \overline{m}) \in I_k(\mathbb{M})$ such that

$$|f(\overline{n}) - f(\overline{m})|_{J_p^{**}} \leq 2Lip(f)k^{\frac{1}{p}} + \varepsilon$$

Proof. We shall prove this statement by induction on $k \in \mathbb{N}$
The proposition is clearly true for $k = 1$.
Assume now that it is true for $k - 1 \geq 1$.
Let $f : G_k(\mathbb{M}) \to J_p^{**}$ be a Lipschitz map and $\varepsilon > 0$.
Let $\eta > 0$ (small enough : to be detailed later).
By a diagonal extraction process and thanks to the weak*-compactness, we can find an infinite subset $\mathbb{M}_1$ of $\mathbb{N}$ such that

$$\forall\ \overline{n} \in G_{k-1}(\mathbb{N}),\ w^* - \lim_{k \in \mathbb{M}_1} f(\overline{n}, k) = g(\overline{n}) \in J_p^{**} \tag{3.2}$$

Then $Lip(g) \leq Lip(f)$, by weak*-lower semicontinuity of $|.|_{J_p^{**}}$.
Denote $g(\overline{n}) = v(\overline{n}) + c_{\overline{n}}\mathbb{1}$ (where $v(\overline{n}) \in J_p$, $c_{\overline{n}} \in \mathbb{R}$ and $\mathbb{1}$ is the constant sequence $(1, 1, 1, ...)$).
By Ramsey's theorem, there exists an infinite subset $\mathbb{M}_2$ of $\mathbb{M}_1$ such that

$$\forall\ \overline{n}, \overline{m} \in G_{k-1}(\mathbb{M}_2) \quad |c_{\overline{n}} - c_{\overline{m}}| = |(v(\overline{n}) - v(\overline{m})) - (g(\overline{n}) - g(\overline{m}))|_{J_p^{**}} < \eta. \tag{3.3}$$

For $\overline{n}, \overline{m} \in G_{k-1}(\mathbb{M}_2)$ and $t, l \in \mathbb{M}_2$, set

$$u_{\overline{n},\overline{m},t,l} = f(\overline{n}, t) - g(\overline{n}) + g(\overline{m}) - f(\overline{m}, l).$$

Using (3.2) and Corollary 2.4 we deduce that there exists $l_0 \in \mathbb{N}$ such that for all $t, l \in \mathbb{M}_2 \cap [l_0,\ +\infty[$:

$$|v(\overline{n}) - v(\overline{m}) + u_{\overline{n},\overline{m},t,l}|^p \leq |v(\overline{n}) - v(\overline{m})|^p + |u_{\overline{n},\overline{m},t,l}|^p + \eta.$$

Then it follows from (3.3) that for all $t, l \in \mathbb{M}_2 \cap [l_0,\ +\infty[$:

$$|f(\overline{n},t) - f(\overline{m},l)|^p_{J_p^{**}} \leq |u_{\overline{n},\overline{m},t,l}|^p_{J_p^{**}} + (|v(\overline{n}) - v(\overline{m})|_{J_p} + \eta)^p + \eta.$$

Moreover : $f(\overline{n},t) - g(\overline{n}) = w^* - \lim_i (f(\overline{n},t) - f(\overline{n},i))$.
Therefore, by weak*-lower semicontinuity of $|.|_{J_p^{**}}$: $|f(\overline{n},t) - g(\overline{n})|_{J_p^{**}} \leq Lip(f)$.
Likewise : $|f(\overline{m},l) - g(\overline{m})|_{J_p^{**}} \leq Lip(f)$.
Then, we deduce the following inequality : $|u_{\overline{n},\overline{m},t,l}|^p_{J_p^{**}} \leq 2^p Lip(f)^p$.
On the other hand, it follows from our induction hypothesis that:

$$\exists\ (\overline{n},\ \overline{m}) \in\ (I_{k-1}(\mathbb{M}_2))^2,\ |g(\overline{n}) - g(\overline{m})|_{J_p^{**}} \leq 2Lip(f)(k-1)^{\frac{1}{p}} + \eta.$$

Then, for $t,\ l \in \mathbb{M}_2 \cap [l_0,\ +\infty[$ such that $m_{k-1} < t < l$, we have $((\overline{n},t),(\overline{m},l)) \in I_k(\mathbb{M}_2)$, and

$$|f(\overline{n},t) - f(\overline{m},l)|^p_{J_p^{**}} \leq 2^p Lip(f)^p + (2Lip(f)(k-1)^{\frac{1}{p}} + 2\eta)^p + \eta.$$

So :

$$|f(\overline{n},t) - f(\overline{m},l)|^p_{J_p^{**}} \leq 2^p Lip(f)^p k + \varphi(\eta),\ \text{with } \varphi(\eta) \underset{\eta\to 0}{\to} 0.$$

Thus, if η was chosen small enough :

$$|f(\overline{n},t) - f(\overline{m},l)|_{J_p^{**}} \leq 2Lip(f)k^{1/p} + \varepsilon.$$

This finishes our inductive proof. □

Corollary 3.6. *Let $1 < q < p < \infty$.*
Then J_q does not coarse Lipschitz embed into J_p.

Proof. Suppose that $g\ : J_q \to J_p$ is a map such that there exist θ, A and B real positive numbers such that :

$$\forall x, y \in J_q\ \ ||x-y||_{J_q} \geq \theta \Rightarrow A||x-y||_{J_q} \leq |g(x) - g(y)|_{J_p} \leq B||x-y||_{J_q}.$$

Let us rescale by defining $f(v) = A\theta^{-1}g(\theta v)$, for $v \in J_q$. We have that there exists $C \geq 1$ such that

$$\forall x, y \in J_q, ||x-y||_{J_q} \geq 1 \Rightarrow ||x-y||_{J_q} \leq |f(x) - f(y)|_{J_p} \leq C||x-y||_{J_q}. \tag{3.4}$$

We still denote $(e_n)_{n=1}^{\infty}$ the canonical basis of J_q.
Consider the map $\varphi\ : G_k(\mathbb{N}) \to (J_q, \|\ \|_{J_q})$ defined by $\varphi(\overline{n}) = e_{n_1} + ... + e_{n_k}$.
Note that $Lip(\varphi) \leq 4$. Since $\|\varphi(\overline{n}) - \varphi(\overline{m})\|_{J_q} \geq 1$ whenever $\overline{n} \neq \overline{m}$, we have that $Lip(f \circ \varphi) \leq 4C$, where $f \circ \varphi$ is considered as a map from $G_k(\mathbb{N})$ to $(J_p, |\ |_{J_p})$. It then follows from Proposition 3.5 that there exist $(\overline{n},\overline{m}) \in I_k(\mathbb{N})$ such that:

$$|(f \circ \varphi)(\overline{n}) - (f \circ \varphi)(\overline{m})|_{J_p} \leq 9Ck^{\frac{1}{p}}.$$

On the other hand, since $(\overline{n},\overline{m}) \in I_k(\mathbb{N})$, we have that

$$||\varphi(\overline{n}) - \varphi(\overline{m})||_{J_q} \geq 2(2k-1)^{\frac{1}{q}} \geq 2k^{\frac{1}{q}}.$$

This is in contradiction with (3.4), for k large enough.
Therefore, there is no coarse Lipschitz embedding from J_q into J_p. □

Corollary 3.7. *Let $1 < p < q < \infty$, and $r > 1$ such that $r \notin \{p,\ q\}$.*
Then J_r does not coarse Lipschitz embed into $J_p \oplus J_q$.

Proof. When $r > q$, the argument is based on a midpoint technique like in the proof of Corollary 3.3.

If $r < p$, we mimic the proof of Corollary 3.6.

So we assume, as we may, that $1 < p < r < q < \infty$ and $f\ : J_r \to J_p \oplus_\infty J_q$ is a map such that there exists $C \geq 1$ such that

$$\forall\ x,\ y \in J_r\ \ |x-y|_{J_r} \geq 1 \Rightarrow |x-y|_{J_r} \leq ||f(x)-f(y)|| \leq C|x-y|_{J_r}. \tag{3.5}$$

We follow the proof in [8] and write $f = (g,h)$. We still denote $(e_n)_{n=1}^\infty$ the canonical basis of J_r. We fix $k \in \mathbb{N}$ and $\varepsilon > 0$. We recall that

$$\exists\ \gamma > 0,\ \forall\ x \in J_r,\ \gamma||x||_{J_r} \leq |x|_{J_r} \leq ||x||_{J_r}.$$

We start by applying the midpoint technique to the coarse Lipschitz map g and deduce from Proposition 3.2 that there exist $\theta > \gamma^{-1}(2k)^{1/r}$, $u \in J_r$, $N \in \mathbb{N}$ and K a compact subset of J_p such that :

$$g(u + \theta B_{(E_N,|.|_{J_r})}) \subset K + \varepsilon\theta B_{(J_p,||.||_{J_p})}. \tag{3.6}$$

Let $\mathbb{M} = \{n \in \mathbb{N},\ n > N\}$ and $\varphi\ : G_k(\mathbb{M}) \mapsto J_r$ be defined as follows

$$\forall\ \overline{n} = (n_1,...,n_k) \in G_k(\mathbb{M}),\ \varphi(\overline{n}) = u + \theta(2k)^{-\frac{1}{r}}(e_{n_1} + ... + e_{n_k}).$$

Then $\varphi(\overline{n}) \in u + \theta B_{(E_N,|.|_{J_r})}$ for all $\overline{n} \in G_k(\mathbb{M})$.
And, from (3.6) we deduce that $(g \circ \varphi)(G_k(\mathbb{M})) \subset K + \varepsilon\theta B_{(J_p,||.||_{J_p})}$. Thus, by Ramsey's theorem, there is an infinite subset $\mathbb{M}'$ of $\mathbb{M}$ such that

$$\mathrm{diam}_{\|\ \|_{J_p}}(g \circ \varphi)(G_k(\mathbb{M}')) \leq 3\varepsilon\theta. \tag{3.7}$$

Since for $\overline{n} \neq \overline{m}$, we have

$$|\varphi(\overline{n}) - \varphi(\overline{m})|_{J_r} \geq \gamma||\varphi(\overline{n}) - \varphi(\overline{m})||_{J_r} \geq \gamma\theta(2k)^{-\frac{1}{r}} > 1,$$

it follows from (3.5) that

$$\forall \overline{n},\overline{m} \in G_k(\mathbb{M})\ \ ||h\circ\varphi(\overline{n}) - h\circ\varphi(\overline{m})||_{J_q} \leq C|\varphi(\overline{n}) - \varphi(\overline{m})|_{J_r} \leq C||\varphi(\overline{n}) - \varphi(\overline{m})||_{J_r}.$$

Note now that for all $\overline{n},\overline{m} \in G_k(\mathbb{N})$,

$$||(e_{n_1} + ... + e_{n_k}) - (e_{m_1} + ... + e_{m_k})||_{J_q} \leq \sum_{n_i \neq m_i} ||e_{n_i} - e_{m_i}||_{J_q} \leq 2d(\overline{n},\overline{m}).$$

Since moreover $|\ |_{J_q} \leq \|\ \|_{J_q}$, $Lip(h\circ\varphi) \leq 2C\theta(2k)^{-\frac{1}{r}}$, when $h\circ\varphi$ is considered as a map from $G_k(\mathbb{M}')$ to $(J_q, |\ |_{J_q})$. Thus, we can apply Proposition 3.5 to obtain:

$$\exists\ (\overline{n},\ \overline{m}) \in I_k(\mathbb{M}'),\ |h\circ\varphi(\overline{n}) - h\circ\varphi(\overline{m})|_{J_q} \leq 5C\theta(2k)^{-\frac{1}{r}}k^{\frac{1}{q}}.$$

Then, if k was chosen large enough, we have:

$$\exists\ (\overline{n},\ \overline{m}) \in I_k(\mathbb{M}'),\ |h \circ \varphi(\overline{n}) - h \circ \varphi(\overline{m})|_{J_q} \leq \varepsilon\theta.$$

This, combined with (3.7) implies that

$$\exists\ (\overline{n},\ \overline{m}) \in I_k(\mathbb{M}')\ \ \|f \circ \varphi(\overline{n}) - f \circ \varphi(\overline{m})\| \leq 3\varepsilon\theta.$$

But,

$$\forall (\overline{n},\ \overline{m}) \in I_k(\mathbb{M}')\ \ |\varphi(\overline{n}) - \varphi(\overline{m})|_{J_r} \geq \gamma \|\varphi(\overline{n}) - \varphi(\overline{m})\|_{J_r} \geq \gamma\theta.$$

If ε was initially chosen such that $\varepsilon < \frac{\gamma}{3}$, this yields a contradiction with (3.5), which concludes our proof. □

Remark. This result can be easily extended as follows. Assume $r \in (1, \infty) \setminus \{p_1,\ ...,\ p_n\}$ where $1 < p_1 < p_2 < ... < p_n < \infty$, then J_r does not coarse Lipschitz embed into $J_{p_1} \oplus ... \oplus J_{p_n}$.

Aknowledgment

The author is grateful to Gilles Lancien for sharing his knowledge and experience, and for helpful discussions that contributed to shape this paper.

References

[1] Y. Benyamini and J. Lindenstrauss, Geometric nonlinear functionan analysis, *A.M.S. Colloquium publications*, vol 48, American Mathematical Society, Providence, RI, (2000).

[2] J. Bourgain, Remarks on the extension of Lipschitz maps defined on discrete sets and uniform homeomorphisms, Geometrical aspects of functional analysis (1985/86), *Lecture Notes in Math., 1267*, Springer Berlin, (1987), 157–167.

[3] E. Gorelik, The uniform nonequivalence of L_p and l_p, *Israel J. Math*, **87**, (1994), 1–8.

[4] W.T. Gowers, Ramsey methods in Banach spaces, *Handbook of the geometry of Banach spaces*, vol 2, North-Holland, Amsterdam, (2003), 10711097.

[5] R.C. James, A somewhat relexive Banach space with non separable dual, *Bull. A.M.S. 80*, (1974), 738–743.

[6] W.B. Johnson, J. Lindenstrauss and G. Schechtman, Banach spaces determined by their uniform structures, *Geom. Funct. Anal.*, **6**, (1996), 430–470.

[7] N.J. Kalton, Coarse and uniform embeddings into reflexive spaces, *Quart. J. Math. (Oxford)*, **58**(1), (2007), 393–414.

[8] N.J. Kalton, N.L. Randrianarivony, The coarse Lipschitz structure of $\ell_p \oplus \ell_q$, *Math. Ann.*, **341**, (2008), 223–237.

[9] G. Lancien, A short course on non linear geometry of Banach spaces. *Topics in Functional and Harmonic Analysis, Theta Series in Advanced Mathematics*, (2012), 77–102.

[10] G. Lancien, Réflexivité et normes duales possédant la propriété uniforme de Kadec-Klee, *Publ. Math. de la Fac. des sciences de Besançon, Fascicule 14*, (1993-94), 69–74.

[11] G. Pisier, The dual J^* of the James Space has cotype 2 and the Gordon-Lewis property, *Proc. Cambridge Phil.*, **103**, (1988), 323–331.

Université de Franche-Comté, Laboratoire de Mathématiques UMR 6623, 16 route de Gray, 25030 Besançon Cedex, FRANCE.

E-mail address: francois.netillard@univ-fcomte.fr